

MINDY S. GIL, DMD, DMSc DIPLOMATE OF AMERICAN BOARD OF PERIODONTOLOGY

1635 GA-34 Suite A, Newnan, GA 30265 p.770.252.0029 info@newnanperio.com f. 770.252.0091 www.newnanperio.com

PREOPERATIVE INSTRUCTIONS FOR I.V. SEDATION

Please AVOID	 Give blood for 5 days before surgery Drink alcohol 24 hours before the surgery Eat or drink anything for 8 hours (food or water) Wear contact lens, jewelry, perfumes, lotions, nail polish
Please DO	 Take all morning medications with a small sip of water (including blood pressure medications) Drink tons of water the day before the surgery Take your antibiotics starting one day PRIOR Arrange for a ride to and from your appointment No ubers, taxis, lyft Arrange for your ride to stay within 10 min radius
Please know	 We may use the following medications during IV sedation: benzodiazepines, narcotics, steroid, NSAIDS, benedryl Some of these medications are category X for pregnancy. If there is any chance you could be pregnant, please let us know.
Please tell us again:	 Antiplatelets you are on: Allergies: A1C for Diabetes (take morning level and let us know): Are you or Are you trying to get pregnant: Yes No

POST INSTRUCTIONS FOR I.V. SEDATION

For the	Drive or operate any machinery
remainder of	Drink alcohol
the day, Please	 Make any important family or legal decisions
DO NÓT	 Do not plan on returning to work or have any other
	appointments the same day as sedation

I have read and understand the preoperative & postoperative instructions for I.V
sedation, and I agree to comply with these instructions.

Signed	Date:	
Witnessed		